



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$r^2 \left[\frac{211}{960} \pi^3 + \frac{207}{360} \pi - .9724 \right] = 160,$$

$$r^2 [6.8148 + 1.8064 - .9726] = 160, \quad r^2 (7.6486) = 160, \quad r^2 = 20.91, \quad r = 4.57.$$

$$2\pi r = 28.714 + \text{the circumference of the wall.}$$

II. Solution by G. B. M. ZERR, A. M., Ph. D., Vice President and Professor of Mathematics in the Texarkana College, Texarkana, Arkansas.

Let ASD , in Fig. above, be the given wall. S the point where the mules are fastened, a =radius of wall, $\varphi = \angle KCL$, $PL = \rho$, =radius of curvature of involute KPF , $\theta = \angle ASC$.

We now have the three areas to find:— (1) area $SHABDKS$, (2) the the two equal involute areas SHE and SKF , (3) area semi-circle EGF . $SA = SE = \frac{1}{2}\pi a$.

$$\therefore \text{Area semi-circle } EGE = \frac{1}{8} \pi^3 a^2 \dots (1).$$

Area of an element between two consecutive radii of curvature is $dA = \frac{1}{2} \rho^2 d\varphi = \frac{1}{2} a^2 \varphi^2 d\varphi$, since $\rho = a\varphi$.

$$\therefore \text{Area } (SHE + SKF) = a^2 \int_0^{\frac{\pi}{2}} \varphi^2 d\varphi = \frac{1}{24} \pi^3 a^2 \dots (2).$$

Area common to both circles = $a^2 (\pi + 2\theta \cos 2\theta - \sin 2\theta)$, but $2a \cos \theta = \frac{1}{2} \pi a$, $\therefore \cos \theta = \frac{1}{4} \pi$.

\therefore Area common to both circles

$$= a^2 \left\{ \pi + \frac{1}{4} (\pi^2 - 8) \cos^{-1} \frac{\pi}{4} - \frac{\pi}{8} \sqrt{16 - \pi^2} \right\} \dots (3).$$

$$\therefore a^2 \left\{ \frac{1}{8} \pi^3 + \pi + \frac{1}{4} (\pi^2 - 8) \cos^{-1} \frac{\pi}{4} - \frac{\pi}{8} \sqrt{16 - \pi^2} \right\} = 160 \text{ sq. rods.}$$

$$\therefore a^2 \left(7.337 + .4674 \cos^{-1} \frac{\pi}{4} \right) = 160 \text{ sq. rods.}$$

$$\therefore 7.64896 a^2 = 160, \quad a = 4.5736 \text{ rds.}$$

$$2\pi a = 28.7368 \text{ rods, = circumference required.}$$

Also solved by A. H. BELL and F. P. MATZ.

PROBLEMS.

45. Proposed by Dr. GEORGE LILLEY, Portland, Oregon.

A fly starts from a point in the circumference of a table, 3 feet in diameter, and travels uniformly along the diameter to a point in the circumference of the table directly opposite the starting point. The table moves uniformly to the right about a center axis in such manner that it makes one complete revolution while the fly passes over its diameter. Find the absolute path described by the fly and the ratio of rates of movement of the table and the fly.